MENTAL A THE Global challenges

Special Issue -

MHGC Proceeding 2020

Kenny Music Performance Anxiety Inventory (K-MPAI): Exploratory Factor Analysis of the Ukrainian version

Olena Ksondzyk

Institute of Social and Political Psychology of National Academy of Educational Science, Kyiv, Ukraine

Abstract

Introduction: Music performance anxiety (MPA) is one of the most common psychological problems among musicians, regardless of their age, gender or level of stage experience. Since empirical studies of this subject are just emerging in Ukraine, there is a lack of psychometrically valid instruments for measuring it. Many specific instruments are available to evaluate MPA in English, but they have to be adapted for the Ukrainian population. One of such significant psychodiagnostic tools is the Kenny Music Performance Anxiety Inventory (K-MPAI) used for different cultural contexts.

Purpose: The aim of this research is to study the factor structure of the Ukrainian version of K-MPAI.

Methodology: In order to assess the K-MPAl's linguistic and conceptual equivalence, the questionnaire was translated using blind back-translation method. Thereafter, the sample of 252 professional musicians (aged 19–66, M=38, SD=11.24; 59% women and 41% men) completed the K-MPAI.

Results: An exploratory factor analysis with principal axis factoring and oblimin rotation method was performed based on the K-MPAI items. The optimal implementation of parallel analysis revealed three factors that explain 44.99 % of variance; they are named "proximal performance concerns", "early relationship context", and "psychological vulnerability". The internal consistency of the Ukrainian version of K-MPAI presents excellent value with Cronbach's alpha of 0.871 and high temporal stability (r = .84; p<0.001).

Discussion & Conclusions: These findings demonstrate evidence of construct validity and reliability of the Ukrainian version of K-MPAI and partially support the theoretical model that became the basis for the development of the original measure. This questionnaire can be used as a valid tool to assess MPA in Ukrainian scientific research.

Keywords

anxiety, performance, musicians, adaptation, test validity, factor structure

Address for correspondence:

Olena Ksondzyk, Laboratory of social psychology of personality, Institute of Social and Political Psychology, NAES of Ukraine, e-mail: olenaksondzyk@gmail.com
This work is licensed under a Creative Commons Attribution-

© (§ (§)

©Copyright: Ksondzyk, 2020

Licensee NDSAN (MFC-Coordinator of the NDSAN), Italy

NonCommercial 4.0 International License (CC BY-NC 4.0).

DOI: http://doi.org/10.32437/mhgcj.v4i2.87

Submitted for publication: 23

May 2020

Received: 23 May 2020 Accepted for publication: 30

September 2020

https://mhgcj.org



MENTAL A THE Global challenges

Special Issue -

MHGC Proceeding 2020

Introduction

Music performance anxiety (MPA) is a particular state of stressful and persistent apprehension related to music performance in front of an audience (Kenny, 2011; Salmon, 1990; Spahn et al., 2016). The core of this anxiety is fear of professional failure, negative evaluation by audience, and possible consequences, though its optimal level is beneficial for the quality of performance. An excessive degree and repeated experiences of performance anxiety can cause increased tension, loss of adequate self-esteem and deterioration of self-efficacy.

MPA is manifested through affective, cognitive, somatic, and behavioural symptoms along a broad continuum of varying severity – from adaptive responses to pathological forms (Kenny, 2011; Spahn et al., 2016). Considering all diagnostic criteria, MPA is classified as a "performance only" subtype of social anxiety disorder in DSM-V (APA, 2013).

Despite the highly developed music education in Ukraine, the empirical studies of music performance anxiety have not been conducted due to several reasons. professional musicians tend to stigmatize and silence this problem, avoiding psychological help. The second significant obstacle to obtaining reliable data is the lack of valid special psychodiagnostic tools for its assessment. In this context, the Kenny Music Performance Anxiety Inventory (K-MPAI) deserves special attention. Based on Barlow's emotion theory of the etiology of anxiety and its disorders (Barlow, 2002), this questionnaire integrates cognitive, emotional, and physiological aspects of MPA taking into account its etiological complexity. The K-MPAI demonstrates excellent psychometric properties (Barbar et al., 2015; Kenny, 2004; Zarza-Alzugaray et al., 2016).

Purpose

The purpose of this research is to study the internal factor structure of K-MPAI based on a sample of the Ukrainian population.

Methodology

Participants

The sample consists of 252 adult musicians from various regions of Ukraine (Lviv, Ivano-Frankivsk, Kyiv, Ternopil, Kharkiv), who work in state institutions of music education (music academies) and philharmonic societies (aged

19-66, M = 38, SD = 11.24; 59% women and 41% men). Ethical approval was obtained from the Laboratory of the Social Psychology of Personality at the Institute of Political and Social Psychology in Kyiv. Thereafter, data have been collected during a concert season that represents a typical schedule of rehearsals and performances. The musicians were guaranteed voluntary participation and anonymity. The criteria for inclusion in the study were age over 18 years old and frequent participation in public musical performances, while the exclusion criterion was the incorrect filling out of the form. Characteristics related to the participants' musical genre and instrument category representation are presented in Table 1.

Table 1: Sample Characteristics (N=252)

Main music genre	N	Percentage (%)
Classical	185	73%
Рор	35	14%
Folk	19	8%
Jazz	7	3%
Rock	6	2%
Instrument		
category	N	Percentage (%)
	N 106	Percentage (%) 42%
category		- , ,
category Piano	106	42%
category Piano Voice	106	42% 15%
category Piano Voice Conducting	106 37 33	42% 15% 13%

Adaptation procedure

Initially, the permission to adapt the K-MPAI for the Ukrainian population was obtained from the author Dianna Theodore Kenny, Professor at the University of Sydney, Australia. Then the questionnaire was translated according to the procedures provided by the requirements for cross-cultural adaptation of the methods outlined in international standards (Bartram, D., & Hambleton, R. K., 2016). The following steps were performed:

- Ukrainian version of the original K-MPAI was prepared.
- Reverse translation into the original language was made by another bilingual translator.

https://mhgcj.org





MHGC Proceeding 2020

- Third specialized translation back into Ukrainian was made.
- The best version was chosen (taking into consideration the following criteria: clarity (intelligibility), conceptual, content, semantic and cultural equivalence). Following the translation procedure, a group of music students (N = 30) completed the questionnaire and stated that all statements were clear.
- Translations were compared, and the most reliable translation was selected.

A professional translator, native Ukrainian and English speakers, as well as several psychologists with clinical experience and previous expertise in adapting measures were involved in the translation of the questionnaire. The final version of the inventory was sent to the Author for approval and then recommended for further adaptation procedures.

Instruments

Kenny Music Performance Anxiety Inventory (K-MPAI) (Kenny et al., 2004) is the scale that includes 26 items implying cognitive, behavioural, and somatic characteristics of anxiety in the context of music performance. It into account Barlow's theoretical components (Barlow, 2002) such as evocation of anxious propositions (e.g., uncontrollability, unpredictability, negative affect, situational cues); attentional shift (e.g., task or self-evaluative focus, fear of negative evaluation); physiological arousal; and memory bias. There have been two published versions of the K-MPAI for adult musicians: the 26-item original version (Kenny et 2004) and the modified 40-item questionnaire (Kenny, 2009). In the current study, we choose the initial version, which is essential for screening.

Data analysis

- The sample's clinical and socialdemographic data are characterized using descriptive statistics.
- The exploratory factorial analysis is carried out to verify the structural validity of the K-MPAI.
- Assessment of the internal consistency of statements is identified based on Cronbach's alpha;
- Temporal stability is measured using the testretest procedure;

Data analysis is performed using SPSS Statistics 20

Results

Descriptive statistics

Descriptive statistics for K-MPAI are presented in Table 2.

Table 2. Mean and standard deviation of the K-MPAI (N = 252).

	Inventor V	Min	Max	M (SD)	Male M (SD)	Female M (SD)
F	K-MPAI	18	124	66 (24)	63(24)	67 (24)

There is no significant gender difference, but women are scored slightly higher. The mean total score on the K-MPAI is higher than the mean score reported by Kenny (2004; N=32, M=54.21, SD=34.21, Minimum = 3, Maximum = 111), but lower than the mean score reported by Zarza-Alzugaray et al. (2016; N=275, M=88.87, SD=21.24).

EFA

The factor structure of the Ukrainian version of the K-MPAI was assessed using a principal axis factoring with oblimin rotation, which is preferable when a correlation between factors exist (Tabachnick and Fidell, 2013). An appropriate Kaiser-Meyer-Olkin (KMO) measure of sampling adequacy and a statistically significant Bartlett's test of sphericity were obtained, thus justifying the use of the factor model (Table 3)

Table 3. KMO and Bartlett's test.

Kaiser-Meyer-Olki Sampling Ad	.892	
Bartlett's Test of Sphericity	Approx. Chi- Square	2515.272
	df	325
	Sig.	< 0.001

We have identified seven factors with eigenvalues greater than Kaiser's criterion of 1. They explain 62.79 % of variance. Factor 1 contributes to more than 30.67 % of the variance, whereas the other factors are less than 10% each. Since Kaiser's method tend to overestimate the number of components (Tabachnick & Fidell, 2013), we apply a parallel analysis with the syntax provided by O'Connor (2000). We have obtained a solution of three factors, which explain 44.99% of variance. We have chosen a cutoff .416, because this best interprets the factor structure. The items which do not meet the loading criterion are deleted (2, 3, 8, 26).

The first factor is named "Proximal performance concerns" (11 items: 7, 10, 12, 13,

N-DSA-N

https://mhgcj.org ISSN 2612-2138



MHGC Proceeding 2020

14, 15, 17, 18, 20, 22, 25); the second factor is named "Early relationship context" (3 items: 9, 19, 24), and the third factor is named "Psychological vulnerability" (8 items: 1, 4, 5, 6, 11, 16, 21, 23). The factor loadings on these three factors are presented in Table 4

Table 4. Factor structure of the K-MPAI for the Ukrainian sample

	Factor		
Item	1	2	3
12 During a performance, I find	.814		
myself thinking about whether I'll			
get through it			
13 Thinking about the evaluation,	.773		
I may get interfered with my			
performance			
10 I never know before a concert	.761		
whether I will perform well			
22 I often prepare for a concert	.756		
with a sense of dread and			
impending disaster			
18 I worry that one bad	.675		
performance may ruin my career			
7 Even if I work hard in	.640		
preparation for a performance, I			
am likely to make mistakes			
14 Even in the most stressful	.633		
performance situations, I am			
confident that I will perform well			
25 I worry so much before a	.628		
performance, I cannot sleep			
15 I am often concerned about a	.627		
negative reaction from the			
audience			
17 From early in my music	.524		
studies, I remember being			
anxious about performing			
20 I give up worthwhile	.497		
performance opportunities due to			
anxiety			
19 My parents always listened to		.780	
me			
9 My parents were mostly		.620	
responsive to my needs		4	
24 My parents encouraged me to		.498	
try new things			
5 Excessive worrying is a			.697
characteristic of my family			/05
16 Sometimes I feel anxious for			.695
no particular reason			(70
1 Sometimes I feel depressed			.678
without knowing why			

11 I often feel that I am not worth		.617
much as a person		
6 I often feel that life has not		.592
much to offer me		
23 I often feel that I have nothing		.569
to look forward to		
4 I often find it difficult to work up		.528
the energy to do things		
21 As a child, I often felt sad		.416

Reliability

The internal consistency of the Ukrainian version of K-MPAI presents excellent value with a Cronbach's alpha of 0.871. The α -Cronbach coefficients for factors 1, 2 and 3 are higher for the Ukrainian factor structure than for the original model (Kenny et al., 2004), Spanish (Zarza-Alzugaray et al., 2016), and Brazilian Portuguese (Barbar et al, 2014; Barbar et al, 2015) (Table 5).

Table 5. α-Cronbach coefficients for the Ukrainian version of K-MPAI, compared to the original and other cultural adaptations

Factor	Ukrainia	Origin	Spanis	Brazilian
	n,	al,	h,	Portugues
	N =	N =	N =	e,
	252	32	490	N = 230
"Proximal				
performanc	.900		.868	.82
e concerns"	.900	-	.000	.02
(11 items)				
"Early				
relationship	.663		.568	.57
context" (3	.003	-	.500	.57
items)				
"Psychologi				
cal	.825		.786	.77
vulnerability"		_		.//
(8 items)				
Total	.871	.944	.866	.82

Temporal stability was measured using the test-retest procedure. The results obtained for 30 participants showed a high temporal stability of the K-MPAI results (r = .84; p<0,001), which suggests good reliability of the K-MPAI as a scale

Discussion & Conclusion

This study aims to adapt to the Ukrainian version of K-MPAI. This measure is chosen among other anxiety evaluation tools since it is designed within D. H. Barlow's theoretical model of emotional disorders, and is adapted by Dianna Kenny for assessing the development and maintenance of MPA (Barlow, 2000; Kenny, 2011).

Adhering to this aim, we provided a double-translation of the 26-item English-language

https://mhgcj.org ISSN 2612-2138





MHGC Proceeding 2020

parent form and conducted a pilot-study to confirm the face validity of the Ukrainian version of the scale, which did not undergo any modifications, compared to the initial version.

Following the EFA procedure using principal axis factoring with oblimin rotation, we obtained three factors related to certain aspects of Barlow's theoretical model. The first factor "Proximal performance concerns" (11 items: 7, 10,12, 13, 14, 15, 17, 18, 20, 22, 25, Cronbach's $\alpha = .900$) reveals specific psychological vulnerability of musicians; the second factor "Early relationship context" (3 items: 9, 19, 24, Cronbach's $\alpha = .663$) demonstrates the features of early family context, and the third factor "Psychological vulnerability" (8 items: 1, 4, 5, 6, 11, 16, 21, 23, Cronbach's $\alpha = .825$) identifies generalized psychological vulnerability. Some items related to the original K-MPAI, such as trust in others (2.8), controllability (3), and reliability of memory (26), demonstrated low factor loading and were not taken into consideration. By its content, the factor model is close to the Spanish version (Zarza-Alzugaray at al., 2016) and Brazilian Portuguese version (Barbar at al., 2015).

The Ukrainian version of K-MPAI presents a high internal consistency with Cronbach's alpha value of .871. High reliability of the K-MPAI as a measure is confirmed by the test-retest procedure that reveals high temporal stability of the K-MPAI results (r = .84; p < 0.001).

Certain limitations of the current study should be noted. First, the sample is composed of adult professional musicians, and most of them are representative of the classical music genre (73%). However, it is important to investigate the manifestations of MPA in samples of other ages (university students in particular) and other musical genres (e.g., folk, jazz, rock, pop). Second in this study, only exploratory factor analysis was performed on the K-MPAI items, but some additional statistical procedures (e.g., CFA) can be carried out to test the resulting three-dimensional model. Future studies could further investigate the convergent and divergent validity of the K-MPAI on the Ukrainian musicians.

The Ukrainian version of K-MPAI has acceptable psychometric properties. Therefore, the measure may be used by specialists as an assessment tool for screening MPA. Since this is the first study to investigate the factorial structure of the K-MPAI in the Ukrainian population, this paper may be useful for further research of MPA and evaluation of treatment efficacy.

Acknowledgment

I would like to express my gratitude to Professor Dianna Kenny for granting the authorization to use the K-MPAI in the current study. I also would like to thank all the musicians, students, and administrative staff of music academies for their participation in the study.

Conflict of interest

The author declares no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

References

American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders (5th ed.). Washington, D C: Author.

Barbar A.E.M., Crippa JAS, Osório F.L. (2014)
Performance anxiety in Brazilian musicians:
Prevalence and association with
psychopathology indicators. Journal of
Affective Disorders. 152(154), 381-6. DOI:
10.1016/j.jad.2013.09.041

Barbar, A. E. M., Souza, J. A. D., & Osório, F. D. L. (2015). Exploratory factor analysis of Kenny Music Performance Anxiety Inventory (K-MPAI) in a Brazilian musician sample. Archives of Clinical Psychiatry (São Paulo), 42(5), 113–116. DOI:10.1590/0101-60830000000060

Barlow D. H. (2002). Anxiety and its disorders: The nature and treatment of anxiety and panic (2nd ed.). New York: The Guilford Press.

Bartram, D., & Hambleton, R. K. (2016). The ITC guidelines: International standards and guidelines relating to tests and testing. In F. T. L. Leong, D. Bartram, F. M. Cheung, K. F. Geisinger, & D. Iliescu (Eds.), The ITC international handbook of testing and assessment (pp. 35–46). Oxford University Press

Kenny, D. T. (2009). The factor structure of the Revised Kenny Music Performance Anxiety Inventory. In A. Williamon (Ed.), Proceedings of the International Symposium on Performance Science (pp. 37–41). Utrecht, The Netherlands: Association Europenne des Conservatoires.

Kenny, D.T. (2011). The Psychology of Music Performance Anxiety. Oxford: Oxford University Press

Kenny, D. T., Davis, P. J., & Oates, J. (2004). Music performance anxiety and occupational stress amongst opera chorus artists and their relationship with state and trait anxiety and perfectionism. Journal of Anxiety Disorders,

https://mhgcj.org ISSN 2612-2138





MHGC Proceeding 2020

- 18(6), 757-777. DOI: 10.1177/0305735612463950
- O'Connor, B. P. (2000). SPSS and SAS programs for determining the number of components using parallel analysis and Velicer's MAP test. Behavior Research Methods, Instruments, & Computers, 32, 396–402. DOI: 10.3758/bf03200807
- Salmon, P. (1990). A psychological perspective on music performance anxiety: A review of the literature. Medical Problems of Performing Artists. 5, 2–11.
- Spahn, C., Walther, J., & Nusseck, M. (2016). The effectiveness of a multimodal concept of

- audition training for music students in coping with music performance anxiety. Psychology of Music, 44 (4), 893-909. DOI: 10.1177/0305735615597484
- Tabachnick, B. G., & Fidell, L. S. (2013). Using multivariate statistics. Upper Saddle River, NJ:Pearson Education.
- Zarza-Alzugaray, F. J., Orejudo, S., Casanova, O., and Mazas, B. (2016). Kenny Music Performance Anxiety Inventory: Confirmatory factor analysis of the Spanish version. Psychol. Music, 44, 340–352. DOI: 10.1177/0305735614567932

https://mhgcj.org

